

THE MASS OF η_b^*

A. PENIN^{a,b}

^a*Department of Physics, University Of Alberta Edmonton, AB T6G 2J1, Canada*

^b*Institute for Nuclear Research, Russian Academy of Sciences 119899 Moscow, Russia*

Abstract

In this paper we briefly review the advances and problems in the QCD theory of the η_b mass.

1 Introduction

The properties of the Υ mesons, the bottom quark-antiquark spin-one bound states, are measured experimentally with great precision, and recent theoretical analysis of the Υ family based on high-order perturbative calculations resulted in determinations of the bottom-quark mass m_b with unprecedented accuracy.^{1,2} At the same time the spin-zero η_b meson remained elusive despite dedicated experimental searches. Only recently a signal of the η_b has been observed by Babar collaboration in the radiative decays of the excited Υ states.^{3,4} The η_b meson shows up as a peak in the photon energy spectrum of the $\Upsilon \rightarrow \gamma\eta_b$ transitions. Despite considerable background from $\Upsilon \rightarrow \gamma\chi_b$ and $e^+e^- \rightarrow \gamma\Upsilon$ processes, the peak energy can be measured with rather high precision. Together with very high accuracy of the Υ spectroscopy, this allows for the determination of η_b mass $M(\eta_b)$ with only a few MeV error. The analysis of the $\Upsilon(3S)$ decays gives $M(\eta_b) = 9388.9^{+3.1}_{-2.3}(\text{stat}) \pm 2.7(\text{syst}) \text{ MeV}$,³ while $\Upsilon(2S)$ data give $M(\eta_b) = 9392.9^{+4.6}_{-4.8}(\text{stat}) \pm 1.9(\text{syst}) \text{ MeV}$.⁴ Thus an accurate prediction of $M(\eta_b)$ is a big challenge and a test for the QCD theory of heavy quarkonium. Due to a very small experimental uncertainty of the $\Upsilon(1S)$ mass, the problem can be reduced to the calculation of the hyperfine splitting (HFS) $E_{\text{hfs}} = M(\Upsilon(1S)) - M(\eta_b)$. This quantity is very sensitive to α_s and could become a competitive source for the determination of the strong coupling constant. In this paper we briefly review the advances and problems in the QCD theory of bottomonium HFS. We consider only the approaches entirely based on the first principles of QCD, leaving aside numerous semi-phenomenological models.

2 Bottomonium Hyperfine Splitting in QCD

Systematic perturbative analysis of the heavy quarkonium bound states is based on the effective field theory of (potential) nonrelativistic QCD, or (p)NRQCD.^{5,6} A recent major breakthrough in the high-order calculations of the heavy quarkonium properties is related to the use of dimensional regularization⁷ and the threshold expansion⁸ within the effective field theory framework.^{2,9} The bottomonium spectrum has been computed to $\mathcal{O}(m_b\alpha_s^5)$, which includes the $\mathcal{O}(\alpha_s)$ next-to-leading order (NLO) correction to the HFS. The corresponding result for an arbitrary principal quantum number is given in Refs.^{2,10} in a closed analytical form. For the ground state it reads

$$E_{\text{hfs}}^{NLO} = \frac{C_F^4 \alpha_s^4 m_b}{3} \left[1 + \frac{\alpha_s}{\pi} \left(\frac{7C_A}{4} \ln(C_F \alpha_s) - \frac{C_F}{2} + \frac{2\pi^2 - 26}{9} n_f T_F + \frac{3 - 3 \ln 2}{2} T_F \right. \right. \\ \left. \left. + \frac{122 - 11\pi^2}{18} C_A \right) \right] \approx E_{\text{hfs}}^{LO} [1 + \alpha_s (1.67 \ln(\alpha_s) + 0.61)] , \quad (1)$$

*Talk given at the 44th Rencontres de Moriond




Figure 1: HFS of 1S bottomonium as a function of the renormalization scale μ in the LO (dotted line), NLO (dashed line), LL (dot-dashed line), and NLL (solid line) approximations. For the NLL result, the band reflects the errors due to $\alpha_s(M_Z) = 0.118 \pm 0.003$.

where $C_F = (N_c^2 - 1)/(2N_c)$, $C_A = N_c = 3$, $n_l = 4$, and α_s is renormalized in the $\overline{\text{MS}}$ scheme at the scale $\mu = C_F \alpha_s m_b$. A logarithmically enhanced term in Eq. (1) is characteristic to the multiscale dynamics of the nonrelativistic bound states.¹¹ Such terms can be resummed to all orders through the renormalization group analysis of pNRQCD, or the *nonrelativistic renormalization group* (NRG)^{12,13} (see also Ref.¹⁴). The renormalization-group-improved expression for the bottomonium HFS is available to the next-to-leading logarithmic (NLL) approximation, which sums up all the corrections of the form $\alpha_s^n \ln^{n-1} \alpha_s$.¹⁵ The corresponding analytical expression is too lengthy to be presented here. The result of the numerical analysis is given in Fig. 1. The logarithmic expansion shows nice convergence and weak scale dependence at the physical scale of the inverse Bohr radius $\mu \sim \alpha_s m_b$. This suggests a small uncertainty due to uncalculated higher-order terms. At the same time the nonperturbative contribution to the HFS is difficult to estimate. In principle it can be investigated by the method of vacuum condensate expansion.¹⁶ The resulting series, however, does not converge well and suffers from large numerical uncertainties.¹⁷ On the other hand, the nonperturbative contribution is suppressed at least by the second power of the heavy quark velocity $v \sim \alpha_s$. Hence it is beyond the accuracy of the NLL approximation and should be added to the errors. In the charmonium system, where the nonperturbative effects are supposed to be much more important, the NLL approximation gives the central value $M(J/\psi) - M(\eta_c) = 104 \text{ MeV}$,¹⁵ which is in a very good agreement with the experimental value $117.7 \pm 1.3 \text{ MeV}$. This suggests that the nonperturbative contribution to the bottomonium HFS is likely to be small as well. A detailed discussion of the uncertainties of the NLL result can be found in Ref.¹⁵ The final numerical prediction for the bottomonium HFS based on perturbative QCD reads

$$E_{\text{hfs}}^{\text{QCD}} = 39 \pm 11 \text{ (th)}_{-8}^{+9} (\delta\alpha_s) \text{ MeV}, \quad (2)$$

where “th” stands for the errors due to the high-order perturbative corrections and nonperturbative effects, whereas “ $\delta\alpha_s$ ” stands for the uncertainty in $\alpha_s(M_Z) = 0.118 \pm 0.003$.

The problem of proper description of the nonperturbative dynamics of strong interactions at long distance is naturally solved by the lattice simulations of QCD. A systematic analysis of the bottomonium HFS within the unquenched lattice NRQCD predicts¹⁸

$$E_{\text{hfs}}^{\text{lat}} = 61 \pm 14 \text{ MeV}, \quad (3)$$

which has somewhat larger central value than Eq. (2), but agrees with the perturbative result within the error bars.

3 Discussion

The estimate (1) based on the perturbative QCD undershoots the experimentally measured values

$$E_{\text{hfs}}^{\text{exp}} = 71.4_{-3.1}^{+2.3} (\text{stat}) \pm 2.7 (\text{syst}) \text{ MeV},^3 \quad E_{\text{hfs}}^{\text{exp}} = 67.4_{-4.6}^{+4.8} (\text{stat}) \pm 2.0 (\text{syst}) \text{ MeV},^4 \quad (4)$$

by about two standard deviations. This discrepancy is rather unexpected and difficult to explain if one takes into account the very successful perturbative description of the HFS in charmonium. At the same time the prediction of the lattice QCD apparently agrees with the experimental data. This fact, however, should be taken with great care. Indeed, the lattice simulation¹⁸ uses a finite lattice spacing $a \sim (\alpha_s m_b)^{-1}$. It is determined by fitting the bottomonium spectrum, which is mostly sensitive to the soft momentum scale $\alpha_s m_b$. At the same time the HFS gets a significant contribution from the hard momentum scale of the heavy quark mass through the radiative corrections. In the lattice NRQCD framework this contribution should be included into the Wilson coefficient of the spin-flip operator in the effective Hamiltonian, which is neglected in Eq. (3). The one-loop Wilson coefficient contains a large logarithm of the form $\ln(am_b)$. It is in one-to-one correspondence with the logarithmic term of Eq. (1) and results in an additional contribution to the HFS

$$\delta^{\text{hard}} E_{\text{hfs}} = -\frac{\alpha_s}{\pi} \frac{7C_A}{4} \ln(am_b) E_{\text{hfs}} \approx -20 \text{ MeV}, \quad (5)$$

which brings the lattice estimate (3) in a perfect agreement with the perturbative result (1). Thus, no definite conclusion on the accuracy of the lattice QCD predictions for the bottomonium HFS can be made at the moment and further theoretical study is necessary. In particular one has to compute the Wilson coefficient of the spin-flip operator perturbatively in the lattice regularization beyond the logarithmic approximation.

To summarize, with the precise experimental data now at hand, the bottomonium HFS becomes one of the most interesting hadronic systems to apply and to test the QCD theory of strong interactions. A significant discrepancy between the prediction based on perturbative QCD and the experimentally measured HFS is intriguing and requires further analysis.

Acknowledgments

This work is supported by the Alberta Ingenuity foundation and NSERC.

References

1. A. Pineda and F.J. Yndurain, Phys. Rev. D **58**, 094022 (1998); J.H. Kühn, A.A. Penin, and A.A. Pivovarov, Nucl. Phys. **B534**, 356 (1998); A.A. Penin and A.A. Pivovarov, Phys. Lett. B **435**, 413 (1998); Nucl. Phys. **B549**, 217 (1999); K. Melnikov and A. Yelkhovsky, Phys. Rev. D **59**, 114009 (1999); M. Beneke and A. Signer, Phys. Lett. B **471**, 233 (1999); A.H. Hoang, Report No. CERN-TH-2000-227 and hep-ph/0008102; A. Pineda, JHEP **0106**, 022 (2001); J.H. Kühn and M. Steinhauser, Nucl. Phys. **B619**, 588 (2001); **B640**, 415(E) (2002).
2. A.A. Penin and M. Steinhauser, Phys. Lett. B **538**, 335 (2002).
3. B. Aubert *et al.* [BABAR Collaboration], Phys.Rev.Lett. **101**, 071801 (2008).
4. B. Aubert *et al.* [BABAR Collaboration], Report No. BaBar-PUB-08/052, SLAC-PUB-13486, and arXiv:0903.1124 [hep-ex].
5. W.E. Caswell and G.P. Lepage, Phys. Lett. B **167**, 437 (1986); G.T. Bodwin, E. Braaten, and G.P. Lepage, Phys. Rev. D **51**, 1125 (1995); **55**, 5853(E) (1997).
6. A. Pineda and J. Soto, Nucl. Phys. B (Proc. Suppl.) **64**, 428 (1998); N. Brambilla, A. Pineda, J. Soto, and A. Vairo, Nucl. Phys. **B566**, 275 (2000).
7. A. Pineda and J. Soto, Phys. Lett. B **420**, 391 (1998); Phys. Rev. D **59**, 016005 (1999).
8. M. Beneke and V.A. Smirnov, Nucl. Phys. **B522**, 321 (1998); V.A. Smirnov, *Applied Asymptotic Expansions in Momenta and Masses* (Springer-Verlag, Heidelberg, 2001).

9. B.A. Kniehl and A.A. Penin, Nucl. Phys. **B563**, 200 (1999); B.A. Kniehl, A.A. Penin, V.A. Smirnov, and M. Steinhauser, Phys. Rev. D **65**, 091503(R) (2002); Nucl. Phys. **B635**, 357 (2002); Phys. Rev. Lett. **90**, 212001 (2003); **91**, 139903(E) (2003); M. Beneke, Y. Kiyo, and A.A. Penin, Phys. Lett. B **653**, 53 (2007).
10. A.A. Penin, V.A. Smirnov, and M. Steinhauser, Nucl. Phys. **B716**, 303 (2005).
11. N. Brambilla, A. Pineda, J. Soto, and A. Vairo, Phys. Lett. B **470**, 215 (1999); B.A. Kniehl and A.A. Penin, Nucl. Phys. **B577**, 197 (2000).
12. A. Pineda, Phys. Rev. D **65**, 074007 (2002); **66**, 054022 (2002).
13. A.A. Penin, A. Pineda, V.A. Smirnov, and M. Steinhauser, Phys. Lett. B **593**, 124 (2004); Nucl. Phys. **B699**, 183 (2004).
14. M.E. Luke, A.V. Manohar, and I.Z. Rothstein, Phys. Rev. D **61**, 074025 (2000).
15. B.A. Kniehl, A.A. Penin, A. Pineda V.A. Smirnov, and M. Steinhauser, Phys. Rev. Lett. **92**, 242001 (2004).
16. M.B. Voloshin, Nucl. Phys. **B154**, 365 (1979); Yad. Fiz. **36**, 247 (1982) [Sov. J. Nucl. Phys. **36**, 143 (1982)]; H. Leutwyler, Phys. Lett. B **98**, 447 (1981).
17. S. Titard and F.J. Yndurain, Phys. Rev. D **51**, 6348 (1995); A. Pineda, Nucl. Phys. **B494**, 213 (1997).
18. A. Gray *et al.*, Phys.Rev. D **72**, 094507 (2005).